



Moustafa M.Hassan, M.D., Director, FACOG  
 Basil Maghak, M.D., FACOG  
 Mohammed A. Elkousy, M.D., FACOG  
 Marquis Jessie, M.D., FACOG

Accredited with the American Institute of Ultrasound in Medicine  
 and by the Fetal Medicine Foundation

**INFORMED CONSENT- Genetic Amniocentesis**

By signing below, I \_\_\_\_\_ request that Genetic Amniocentesis be performed by Dr. \_\_\_\_\_ for the purpose of prenatal diagnosis of certain birth defects. It has been explained to me by my doctor or by someone my doctor has designated, and I understand that:

- 1-- Amniocentesis is the withdrawal of a small sample (less than 1 oz) of the fluid surrounding the fetus. This fluid is obtained by inserting a needle through the abdominal wall into the uterus (womb). Ultrasound is usually performed to help locate the placenta and fetus. Ultrasound may also detect twins, maybe used to date the pregnancy, and may detect some but not all physical defects in the fetus.
- 2-- Amniocentesis involves a small risk to both the mother and fetus. The most common serious complication is miscarriage; in general, the risk of miscarriages from amniocentesis performed after 15 weeks of pregnancy is less than 1 in 250. Other possible but serious complications include hemorrhage, infection or injury to the fetus. Minor complications, which occur in approximately 1 in 100 woman having amniocentesis, include cramping, vaginal spotting, or slight leakage of amniotic fluid.
- 3-- Any particular attempt to obtain amniotic fluid may be unsuccessful. Occasionally, even if sufficient fluid is obtained, laboratory testing may not be possible or may not yield results. In these cases, the amniocentesis may need to be repeated.
- 4-- The standard laboratory testing preformed on the amniotic fluid sample consists of chromosome analysis, which can usually also detect a high percentage of all open neural tube defects.
- 5-- Testing for other kinds of birth defects will not be performed unless indicated below.
- 6-- Normal test results do not guarantee the birth of a normal child. As in any laboratory test, there is a small possibility of error, and maternal cells may contaminate the sample. In addition, approximately 2---3% of all pregnancies have birth defects which cannot be detected by testing amniotic fluid or by ultrasound examination.
- 7-- Any part of this amniotic fluid sample not used for diagnostic testing maybe stored and used for medical research or education as long as my name or any other identifying information have been removed.

The reason I am having an amniocentesis today is: \_\_\_\_\_

\_\_\_\_\_    \_\_\_/\_\_\_/\_\_\_\_    \_\_\_\_\_    \_\_\_/\_\_\_/\_\_\_\_  
 Patient Signature                      Date                      Witness                      Date

1850 Town Center Pkwy  
 Suite 258  
 Reston, VA 20190-3219  
 (703) 435-1454  
 (703) 435-8630 fax

44035 Riverside Parkway  
 Suite 345  
 Lansdowne, VA 20176-8273  
 (703) 724-4250  
 (703) 724-4218 fax

7915 Lake Manassas Drive  
 Suite 210  
 Gainesville, VA 20155-3260  
 (703) 743-9211  
 (703) 724-1659 fax

3580 Joseph Siewick Drive  
 Suite 304  
 Fairfax, VA 22033-1751  
 (703) 584-7701  
 (703) 724-7693 fax